

Note di Matematica **26**, n. 1, 2006, 69–77.

Simply connected two-step homogeneous nilmanifolds of dimension 5

Szilvia Homolya

*Institute of Mathematics, University of Miskolc,
Hungary*
`mathszil@gold.uni-miskolc.hu`

Oldřich Kowalski

*Faculty of Mathematics and Physics, Charles University,
Czech Republic*
`kowalski@karlin.mff.cuni.cz`

Received: 11/3/2005; accepted: 11/3/2005.

Abstract. The aim of this paper is to classify all simply connected two-step nilpotent Lie groups of dimension 5 equipped with left-invariant metrics (“two-step nilmanifolds”) up to isometry. We also calculate the corresponding isometry groups.

Keywords: nilmanifolds, two-step nilpotent Lie groups, H-type groups

MSC 2000 classification: 22E25, 53C30

1 Introduction

Two-step nilpotent Lie groups endowed with a left-invariant metric, often called two-step homogeneous nilmanifolds are studied intensively in the last twenty years. A special subclass of two-step homogeneous nilmanifolds, the Heisenberg type groups, was introduced and studied by A. Kaplan (cf. [6], [7]) and others. We refer to [1] for a survey about the geometry of generalized Heisenberg groups. The Heisenberg type groups play an important role in geometric analysis, Lie groups and mathematical physics. J. Lauret (cf. [10]) essentially generalized this concept by introducing so-called *modified H-type groups* (See below).

In [9] J. Lauret classified, up to isometry, all homogeneous nilmanifolds of dimension 3 and 4 (not necessarily two-step nilpotent) and computed the corresponding isometry groups. He also studied, as example, the structure of specific 5-dimensional two-step nilmanifolds with 2-dimensional center. His results will be used in the present paper. Our purpose is to classify all simply connected two-step Riemannian nilmanifolds of dimension 5 and to determine their full isometry groups.

2 Two-step nilpotent Lie groups

A connected Riemannian manifold which admits a transitive nilpotent Lie group N of isometries is called a *nilmanifold*. E. Wilson proved in [12] that, when given a homogeneous nilmanifold M , there exists a unique nilpotent Lie subgroup N of $I(M)$ acting simply transitively on M , and N is normal in $I(M)$. Hence the Riemannian manifold M can be identified with the group N equipped with a left-invariant metric $\langle \cdot, \cdot \rangle$. A left-invariant metric $\langle \cdot, \cdot \rangle$ on N determines an inner product $\langle \cdot, \cdot \rangle$ on the corresponding Lie algebra $\mathfrak{n} = T_e N$ and conversely. According to [12], the full group of isometries of $(N, \langle \cdot, \cdot \rangle)$ can be expressed as a semi-direct product

$$I(N, \langle \cdot, \cdot \rangle) = K \ltimes N, \quad (1)$$

where $K = \text{Aut}(\mathfrak{n}) \cap O(\mathfrak{n}, \langle \cdot, \cdot \rangle)$ is the isotropy subgroup at the identity element e and N acts by left translations. K is the full group of automorphisms of the Lie algebra \mathfrak{n} which preserve the inner product $\langle \cdot, \cdot \rangle$. Thus the structure of the full isometry group is completely determined by the isotropy subgroup K . Moreover, if N is simply connected, then the exponential mapping $\exp : \mathfrak{n} \rightarrow N$ is a diffeomorphism. We need not make distinction between automorphisms of \mathfrak{n} and those of N .

A Lie algebra \mathfrak{n} is said to be *two-step nilpotent* if $[\mathfrak{n}, \mathfrak{n}] \neq \{0\}$ but $[[\mathfrak{n}, \mathfrak{n}], \mathfrak{n}] = \{0\}$. In the following we shall work usually with a two-step nilpotent metric Lie algebra $(\mathfrak{n}, \langle \cdot, \cdot \rangle)$. We denote by \mathfrak{z} the center of \mathfrak{n} and by $\mathfrak{a} = \mathfrak{z}^\perp$ the orthogonal complement of \mathfrak{z} . Then we have the orthogonal direct sum decomposition $\mathfrak{n} = \mathfrak{a} \oplus \mathfrak{z}$. We denote by $so(\mathfrak{a})$ the Lie algebra of skew-symmetric endomorphisms of the Euclidean vector subspace $(\mathfrak{a}, \langle \cdot, \cdot \rangle_{\mathfrak{a}})$ of $(\mathfrak{n}, \langle \cdot, \cdot \rangle)$.

1 Definition. Let $(\mathfrak{n}, \langle \cdot, \cdot \rangle)$ be a two-step nilpotent metric Lie algebra, $\mathfrak{n} = \mathfrak{a} \oplus \mathfrak{z}$. For each element Z of \mathfrak{z} , define an endomorphism $j(Z) \in so(\mathfrak{a})$ by

$$\langle j(Z)X, Y \rangle = \langle [X, Y], Z \rangle \text{ for all } X, Y \in \mathfrak{a}. \quad (2)$$

The algebraic properties of $(\mathfrak{n}, \langle \cdot, \cdot \rangle)$ can be expressed in terms of the maps $j(Z)$ with $Z \in \mathfrak{z}$. Indeed, let two metric vector spaces $(\mathfrak{a}, \langle \cdot, \cdot \rangle_{\mathfrak{a}})$ and $(\mathfrak{z}, \langle \cdot, \cdot \rangle_{\mathfrak{z}})$ be given and let $(\mathfrak{n}, \langle \cdot, \cdot \rangle_{\mathfrak{n}})$ be their orthogonal direct sum. Let a linear map $j : \mathfrak{z} \rightarrow so(\mathfrak{a})$ be fixed. Define first a Lie bracket on \mathfrak{a} by the condition $[\mathfrak{a}, \mathfrak{a}] \subset \mathfrak{z}$ and by (2). Then extend this Lie bracket to the whole of \mathfrak{n} by putting $[\mathfrak{n}, \mathfrak{z}] = \{0\}$. Since $[\mathfrak{n}, \mathfrak{n}] = [\mathfrak{a}, \mathfrak{a}] \subset \mathfrak{z}$ and $[\mathfrak{n}, \mathfrak{z}] = \{0\}$, the algebra \mathfrak{n} (and hence the corresponding simply connected group N with the left-invariant metric $\langle \cdot, \cdot \rangle$) is two-step nilpotent. The mappings $\{j(Z) : Z \in \mathfrak{z}\}$ contain the “geometry of $(N, \langle \cdot, \cdot \rangle)$ ” in the sense that the covariant derivative, curvature tensor and Ricci tensor can be formulated entirely using j, \mathfrak{a} and \mathfrak{z} (cf. [2]). If \mathfrak{z} is equal to the commutator of \mathfrak{n} , i.e. $\mathfrak{z} = [\mathfrak{n}, \mathfrak{n}]$, then $j : \mathfrak{z} \rightarrow so(\mathfrak{a})$ is an injective linear map.

Among two-step nilpotent Lie groups with left-invariant metrics, the Heisenberg type Lie groups are of particular significance. A two-step nilpotent Lie group N with a left-invariant metric $\langle \cdot, \cdot \rangle$ is said to be an *Heisenberg type* (H-type) Lie group if $[j(Z)]^2 = -\langle Z, Z \rangle id_{\mathfrak{a}}$ for any $Z \in \mathfrak{z}$.

More generally, $(N, \langle \cdot, \cdot \rangle)$ is called a *modified H-type group* if $[j(Z)]^2 = -h(Z)id_{\mathfrak{a}}$ for any $Z \in \mathfrak{z}$, where $h(Z)$ is a positive definite quadratic form on \mathfrak{z} (see [10]).

2 Example. *The Heisenberg group* is, up to isomorphism, the only two-step nilpotent Lie group with a 1-dimensional center. The $(2n+1)$ -dimensional Heisenberg group H_{2n+1} is the group of all real $(n+2) \times (n+2)$ matrices of the form

$$\begin{pmatrix} 1 & x_1 & x_2 & \cdots & x_n & z \\ 0 & 1 & 0 & \cdots & 0 & y_1 \\ \vdots & 0 & 1 & 0 & \vdots & y_2 \\ \vdots & & \ddots & \ddots & 0 & \vdots \\ \vdots & & & \ddots & 1 & y_n \\ 0 & \cdots & \cdots & \cdots & 0 & 1 \end{pmatrix}$$

for $x_i, y_i, z \in \mathbb{R}, i = 1, \dots, n$. The Lie algebra \mathfrak{h}_{2n+1} of H_{2n+1} is a $(2n+1)$ -dimensional vector space with basis $\{X_1, \dots, X_n, Y_1, \dots, Y_n, Z\}$ and the only nonzero Lie brackets are $[X_i, Y_i] = -[Y_i, X_i] = Z$ for $1 \leq i \leq n$.

The center \mathfrak{z} of \mathfrak{h}_{2n+1} equals $\mathfrak{z} = \text{span}_{\mathbb{R}}\{Z\}$. One determines a left-invariant metric on H_{2n+1} by specifying an inner product $\langle \cdot, \cdot \rangle$ on \mathfrak{h}_{2n+1} .

Now we choose the natural inner product making the basis above an orthonormal basis. Using the equation (2) and the basis $\{X_1, \dots, X_n, Y_1, \dots, Y_n, Z\}$ we obtain that

$$j(Z)X_i = Y_i \text{ and } j(Z)Y_i = -X_i \quad \text{for } 1 \leq i \leq n.$$

It follows that $[j(Z)]^2 = -id_{\mathfrak{a}}$, which means that the Heisenberg Lie group equipped with the corresponding natural left-invariant metric is an H-type Lie group.

3 5-dimensional two-step nilmanifolds

Now we give a classification of 5-dimensional simply connected two-step nilmanifolds up to isometry. This is equivalent to the classification of the corresponding metric Lie algebras. Clearly, the dimension of the center of a 5-dimensional two-step nilpotent Lie algebra is ≤ 3 . We consider separately the cases where the dimension of the center is 1, 2 or 3.

3.1 Metric Lie algebras with 1-dimensional center

Let \mathfrak{h}_5 denotes a 5-dimensional Lie algebra the center \mathfrak{z} of which is one-dimensional. We assume that \mathfrak{h}_5 is equipped with an inner product $\langle \cdot, \cdot \rangle$. (Two-step nilmanifolds with one-dimensional center are usually called Heisenberg manifolds.) Let e_5 be a unit vector in \mathfrak{z} and let \mathfrak{a} be the orthogonal complement of \mathfrak{z} in \mathfrak{h}_5 . We consider a 2-dimensional vector subspace \mathfrak{a}_2 such that $[\mathfrak{a}_2, \mathfrak{a}_2] = \mathfrak{z}$. Since the center of \mathfrak{h}_5 is one-dimensional there is a unique vector subspace \mathfrak{b}_2 of \mathfrak{a} which is complementary to \mathfrak{a}_2 in \mathfrak{a} and commutes with \mathfrak{a}_2 , namely $\mathfrak{b}_2 = \text{Ker}(ad(u)) \cap \text{Ker}(ad(v)) \cap \mathfrak{a}$, where $\{u, v\}$ is any basis of \mathfrak{a}_2 . Let us assume that the complementary subspaces \mathfrak{a}_2 and \mathfrak{b}_2 are not orthogonal. One can see easily that the angle of a variable vector in \mathfrak{a}_2 and of its orthogonal projection on \mathfrak{b}_2 achieves its maximum and minimum values in two mutually perpendicular directions in \mathfrak{a}_2 . This angle can be also constant, then the perpendicular directions can be chosen in arbitrary way. (In such a case, \mathfrak{a}_2 and \mathfrak{b}_2 are called isoclinic.) Then one can choose an orthonormal basis $\{e_1, e_2, e_3, e_4\}$ for \mathfrak{a} such that $\{e_1, e_2\}$ is an orthonormal basis of \mathfrak{a}_2 and $\{f_1 = \cos \alpha e_1 + \sin \alpha e_3, f_2 = \cos \beta e_2 + \sin \beta e_4\}$ forms an orthonormal basis for \mathfrak{b}_2 , where $\alpha, \beta \neq 0$ denote the extremal values of the angles between the two-spaces \mathfrak{a}_2 and \mathfrak{b}_2 . We have

$$[e_1, e_2] = \bar{\lambda} e_5 \quad \text{and} \quad [e_3, e_4] = \bar{\mu} e_5$$

where $\bar{\lambda} \neq 0$ and e_5 is a unit vector of the center. Then we can compute the remaining Lie brackets of the basis elements:

$$\begin{aligned} [e_1, e_3] &= 0, & [e_1, e_4] &= -\bar{\lambda} \cot \beta e_5 \\ [e_2, e_4] &= 0, & [e_2, e_3] &= \bar{\lambda} \cot \alpha e_5. \end{aligned}$$

Now, we can find a new orthonormal basis of the form:

$$\begin{aligned} e'_1 &= \cos t e_1 + \sin t e_3, & e'_3 &= -\sin t e_1 + \cos t e_3, \\ e'_2 &= \cos s e_2 + \sin s e_4, & e'_4 &= -\sin s e_2 + \cos s e_4, \end{aligned}$$

such that $[e'_2, e'_3] = 0$ and $[e'_1, e'_4] = 0$. Such a basis is determined by the solution $\{t, s \in \mathbb{R}\}$ of the equations

$$\begin{aligned} (\bar{\lambda} + \bar{\mu}) \sin(t - s) &= \bar{\lambda} (\cot \beta - \cot \alpha) \cos(t - s), \\ (\bar{\lambda} - \bar{\mu}) \sin(t + s) &= -\bar{\lambda} (\cot \beta + \cot \alpha) \cos(t + s). \end{aligned}$$

Hence there exists an orthonormal basis $\{e_1, e_2, e_3, e_4\}$ of \mathfrak{a} such that

$$[e_1, e_2] = -[e_2, e_1] = \lambda e_5, \quad [e_3, e_4] = -[e_4, e_3] = \mu e_5, \quad (3)$$

$$[e_1, e_3] = [e_1, e_4] = [e_2, e_3] = [e_2, e_4] = 0. \quad (4)$$

Moreover we can assume that $\lambda \geq \mu > 0$. A *metric Heisenberg algebra of type (λ, μ)* is defined as a 5-dimensional metric Lie algebra having an orthonormal basis $\{e_1, e_2, e_3, e_4, e_5\}$ satisfying the commutation relations (3) and (4), where $\lambda \geq \mu > 0$. We will denote it by $\mathfrak{h}_5(\lambda, \mu)$.

3 Proposition. *For any 5-dimensional 2-step nilpotent metric Lie algebra \mathfrak{n} with 1-dimensional center there exist real numbers $\lambda \geq \mu > 0$ such that \mathfrak{n} is isomorphic to the metric Heisenberg algebra $\mathfrak{h}_5(\lambda, \mu)$. The metric Heisenberg algebras $\mathfrak{h}_5(\lambda, \mu)$ and $\mathfrak{h}_5(\lambda', \mu')$ are isometrically isomorphic if and only if $\lambda = \lambda'$ and $\mu = \mu'$.*

PROOF. The first assertion follows from the previous discussion.

Let us consider the metric Heisenberg algebras $\mathfrak{h}_5(\lambda, \mu)$ and $\mathfrak{h}_5(\lambda', \mu')$ given by the orthonormal bases $\{e_1, e_2, e_3, e_4, e_5\}$ and $\{e'_1, e'_2, e'_3, e'_4, e'_5\}$, respectively. A linear isomorphism $\varphi : \mathfrak{h}_5(\lambda, \mu) \rightarrow \mathfrak{h}_5(\lambda', \mu')$ is an isometric Lie algebra isomorphism only if it maps the center $\text{span}(e_5)$ onto the center $\text{span}(e'_5)$ and the orthogonal complement \mathfrak{a} onto the orthogonal complement \mathfrak{a}' . In particular $\varphi(e_5) = \pm e'_5$. Hence the maps $j(e_5)^2 : \mathfrak{a} \rightarrow \mathfrak{a}$ and $j(e'_5)^2 : \mathfrak{a}' \rightarrow \mathfrak{a}'$ are selfadjoint endomorphisms with eigenvalues $-\lambda^2, -\mu^2$ and $-\lambda'^2, -\mu'^2$, respectively. We have the relation

$$\varphi \circ j(e_5)^2 \circ \varphi^{-1} = j(e'_5)^2.$$

Hence any isometric isomorphism $\varphi : \mathfrak{h}_5(\lambda, \mu) \rightarrow \mathfrak{h}_5(\lambda', \mu')$ maps the eigenspaces of $j(e_5)^2$ into the eigenspaces of $j(e'_5)^2$ corresponding to the same eigenvalue. It follows $\lambda = \lambda'$ and $\mu = \mu'$. QED

From the above computations and also from [8] we get the following:

4 Corollary. *Each 5-dimensional Heisenberg group space N corresponding to a metric algebra $\mathfrak{h}_5(\lambda, \mu)$ is a modified H -type group in the sense of J. Lauret and it is naturally reductive. It is an H -type group if and only if $\lambda = \mu$.*

If now $\lambda \neq \mu$, then the group of isometric isomorphisms of $\mathfrak{h}_5(\lambda, \mu)$ can be represented by the group of matrices

$$\left\{ \begin{pmatrix} \varepsilon \cos t & -\sin t & 0 & 0 & 0 \\ \varepsilon \sin t & \cos t & 0 & 0 & 0 \\ 0 & 0 & \varepsilon \cos s & -\sin s & 0 \\ 0 & 0 & \varepsilon \sin s & \cos s & 0 \\ 0 & 0 & 0 & 0 & \varepsilon \end{pmatrix}, \quad \varepsilon = \pm 1, s, t \in \mathbb{R} \right\}.$$

In the case $\lambda = \mu$ an orthogonal transformation φ is an orthogonal automorphism if and only if its restriction to \mathfrak{a} commutes with the complex structure $J : \mathfrak{a} \rightarrow \mathfrak{a}$ determined by

$$J(e_1) = \varepsilon e_2, \quad J(e_2) = -\varepsilon e_1, \quad J(e_3) = \varepsilon e_4, \quad J(e_4) = -\varepsilon e_3$$

and, moreover,

$$\varphi(e_5) = \varepsilon e_5, \text{ where } \varepsilon = \pm 1.$$

Hence one obtains:

5 Proposition. *The group of orthogonal automorphisms of the metric Lie algebra $\mathfrak{h}_5(\lambda, \mu)$ is isomorphic to the group $O(2) \times SO(2)$ for $\lambda \neq \mu$, and it is isomorphic to the group $U(2) \times \mathbb{Z}_2$ for $\lambda = \mu$.*

3.2 Metric Lie algebras with 2-dimensional center

Let \mathfrak{n}_5 denotes a 5-dimensional Lie algebra the center \mathfrak{z} of which is two-dimensional and let N_5 be the corresponding simply connected Lie group. We assume that \mathfrak{n}_5 is equipped with an inner product $\langle \cdot, \cdot \rangle$. Let \mathfrak{a} denote the orthogonal complement of the center \mathfrak{z} in \mathfrak{n}_5 . Since the 3-dimensional vector space \mathfrak{a} is isomorphic to the exterior product $\mathfrak{a} \wedge \mathfrak{a}$, the linear map $[\cdot, \cdot] : \mathfrak{a} \wedge \mathfrak{a} \rightarrow \mathfrak{z}$ has a one-dimensional kernel spanned by a bivector $u \wedge v$. The two-dimensional subalgebra $\mathfrak{a}_2 = \mathbb{R}u + \mathbb{R}v$ is a unique two-dimensional commutative subalgebra in \mathfrak{a} . Let $e_1 \in \mathfrak{a}$ be a unit vector which is orthogonal to \mathfrak{a}_2 and let $\{e'_2, e'_3\}$ be an orthonormal basis for \mathfrak{a}_2 . If we consider a new orthonormal basis of \mathfrak{a}_2 in the form

$$e_2 = \cos t e'_2 + \sin t e'_3, \quad e_3 = -\sin t e'_2 + \cos t e'_3, \quad t \in \mathbb{R},$$

then we have

$$\begin{aligned} \langle [e_1, e_2], [e_1, e_3] \rangle = \\ \frac{1}{2}(\| [e_1, e'_3] \|^2 - \| [e_1, e'_2] \|^2) \sin 2t + \langle [e_1, e'_2], [e_1, e'_3] \rangle \cos 2t. \end{aligned} \quad (5)$$

Clearly, one can find $t \in \mathbb{R}$ such that

$$\langle [e_1, e_2], [e_1, e_3] \rangle = 0 \quad (6)$$

and hence the vectors $[e_1, e_2], [e_1, e_3] \in \mathfrak{z}$ are orthogonal (and we still have $[e_2, e_3] = 0$). Thus there is an orthonormal basis $\{e_4, e_5\}$ of \mathfrak{z} such that

$$[e_1, e_2] = \lambda e_4, \quad [e_1, e_3] = \mu e_5, \quad \lambda \geq \mu > 0. \quad (7)$$

We denote a 5-dimensional metric Lie algebra described above by $\mathfrak{n}_5(\lambda, \mu)$. From the considerations in [9, p. 153], we obtain at once

6 Proposition. *For any 5-dimensional 2-step nilpotent metric Lie algebra \mathfrak{n} having a 2-dimensional center there exist real numbers $\lambda \geq \mu > 0$ such that \mathfrak{n} is isomorphic to the metric algebra $\mathfrak{n}_5(\lambda, \mu)$. Moreover the metric Heisenberg Lie algebras $\mathfrak{n}_5(\lambda, \mu)$ and $\mathfrak{n}_5(\lambda', \mu')$ are isometrically isomorphic if and only if $\lambda = \lambda'$ and $\mu = \mu'$.*

7 Proposition. *The group of orthogonal automorphisms of the metric Lie algebra $\mathfrak{n}_5(\lambda, \mu)$ is isomorphic to the group $\mathbb{Z}_2 \times \mathbb{Z}_2 \times \mathbb{Z}_2$ for $\lambda \neq \mu$, and it is isomorphic to the group $O(2) \times \mathbb{Z}_2$ for $\lambda = \mu$.*

Using (2) and (7) we see that, for any $Z = \alpha e_4 + \beta e_5 \in \mathfrak{z}$, the map $j(Z)$ and its square $j^2(Z)$ can be expressed by

$$\begin{aligned} j(Z) &= \begin{pmatrix} 0 & -\alpha\lambda & -\beta\mu \\ \alpha\lambda & 0 & 0 \\ \beta\mu & 0 & 0 \end{pmatrix}; \\ j^2(Z) &= \begin{pmatrix} -\alpha^2\lambda^2 - \beta^2\mu^2 & 0 & 0 \\ 0 & -\alpha^2\lambda^2 & -\alpha\beta\lambda\mu \\ 0 & -\alpha\beta\lambda\mu & -\beta^2\mu^2 \end{pmatrix}. \end{aligned}$$

We see that the 5-dimensional group spaces corresponding to the metric algebras $\mathfrak{n}_5(\lambda, \mu)$ are not modified H-type groups. From [8] we also see easily that these spaces are never naturally reductive.

3.3 Metric Lie algebras with 3-dimensional center

Let \mathfrak{z} denote the center of a two-step nilpotent 5-dimensional metric Lie algebra \mathfrak{n} , where $\dim\{\mathfrak{z}\} = 3$. Clearly $\dim\{[\mathfrak{n}, \mathfrak{n}]\} = 1$ for the commutator $[\mathfrak{n}, \mathfrak{n}]$ of the Lie algebra \mathfrak{n} . Let \mathfrak{a} denote the orthogonal complement of the center \mathfrak{z} in \mathfrak{n} and let \mathfrak{b} denote the orthogonal complement of $[\mathfrak{n}, \mathfrak{n}]$ in the center \mathfrak{z} . Then $[\mathfrak{n}, \mathfrak{n}] = [\mathfrak{a}, \mathfrak{a}]$ and the subalgebra $\mathfrak{h}_3 = \mathfrak{a} \oplus [\mathfrak{a}, \mathfrak{a}]$ is a 3-dimensional metric Heisenberg algebra. The metric Lie algebra \mathfrak{n} decomposes into the orthogonal direct sum $\mathfrak{n} = \mathfrak{h}_3 \oplus \mathfrak{b}$ of the metric Heisenberg subalgebra \mathfrak{h}_3 and of the abelian metric algebra \mathfrak{b} .

Let $\{e_1, e_2\}$ be an orthonormal basis for \mathfrak{a} and $e_3 \in [\mathfrak{a}, \mathfrak{a}]$ a unit vector such that

$$[e_1, e_2] = -[e_2, e_1] = \lambda e_3 \quad (8)$$

with $\lambda > 0$. Moreover we denote by $\{e_4, e_5\}$ an orthonormal basis for \mathfrak{b} . The corresponding Lie algebra will be denoted by $(\mathfrak{h}_3)(\lambda) \oplus \mathbb{R}^2$.

Using the results from [9, pp. 148–149], or by an easy calculation we get

8 Proposition. *For any 5-dimensional 2-step nilpotent metric Lie algebra \mathfrak{n} having a 3-dimensional center there exist a real number $\lambda > 0$ such that \mathfrak{n} is isomorphic to the metric algebra $\mathfrak{h}_3(\lambda) \oplus \mathbb{R}^2$. The metric Heisenberg Lie algebras $\mathfrak{h}_3(\lambda) \oplus \mathbb{R}^2$ and $\mathfrak{h}_3(\lambda') \oplus \mathbb{R}^2$ are isometrically isomorphic if and only if $\lambda = \lambda'$.*

9 Proposition. *The group of orthogonal automorphisms of the metric Lie algebra $\mathfrak{h}_3 \oplus \mathbb{R}^2$ is isomorphic to the group $O(2) \times O(2)$.*

Now we prove that the 5-dimensional group spaces corresponding to the metric algebras $\mathfrak{h}_3(\lambda) \oplus \mathbb{R}^2$ are not modified H-type groups. Using the formulas (2) and (8), we obtain the following matrix representations for $j(e_3)$, $j(e_4)$ and $j(e_5)$:

$$j(e_3) = \begin{pmatrix} 0 & \lambda \\ -\lambda & 0 \end{pmatrix}; \quad j(e_4) = j(e_5) = \begin{pmatrix} 0 & 0 \\ 0 & 0 \end{pmatrix}.$$

We can see that, for any $Z = \alpha e_3 + \beta e_4 + \gamma e_5 \in \mathfrak{z}$, the map $j^2(Z)$ has the form

$$j^2(Z) = \begin{pmatrix} -\alpha^2 \lambda^2 & 0 \\ 0 & -\alpha^2 \lambda^2 \end{pmatrix}.$$

Hence $-j^2(Z)$ is only positive semidefinite. On the other hand, it is known that all these spaces are naturally reductive.

3.4 Classification up to isometry

We can summarize our results:

10 Theorem. *The simply connected two-step homogeneous nilmanifolds of dimension 5 are, up to isometry,*

$$\begin{aligned} (H_5, \langle \cdot, \cdot \rangle_{\lambda, \mu}) & : \quad \lambda \geq \mu > 0, \\ (N_5, \langle \cdot, \cdot \rangle_{\lambda, \mu}) & : \quad \lambda \geq \mu > 0, \\ (H_3 \times \mathbb{R}^2, \langle \cdot, \cdot \rangle_{\lambda}) & : \quad \lambda > 0. \end{aligned}$$

Furthermore the full isometry groups of the corresponding nilmanifolds are expressed by:

$$\begin{aligned} I(H_5, \langle \cdot, \cdot \rangle_{\lambda, \mu}) & = \begin{cases} (U(2) \times \mathbb{Z}_2) \ltimes H_5, & \text{if } \lambda = \mu, \\ (O(2) \times SO(2)) \ltimes H_5, & \text{if } \lambda \neq \mu, \end{cases} \\ I(N_5, \langle \cdot, \cdot \rangle_{\lambda, \mu}) & = \begin{cases} (O(2) \times \mathbb{Z}_2) \ltimes N_5, & \text{if } \lambda = \mu, \\ (\mathbb{Z}_2 \times \mathbb{Z}_2 \times \mathbb{Z}_2) \ltimes N_5, & \text{if } \lambda \neq \mu, \end{cases} \\ I(H_3 \times \mathbb{R}^2, \langle \cdot, \cdot \rangle_{\lambda}) & = (O(2) \times O(2)) \ltimes (H_3 \times \mathbb{R}^2). \end{aligned}$$

PROOF. The result follows at once from the one-to-one correspondence between simply connected two-step nilmanifolds and metric Lie algebras. Here we just use Propositions 3-9. **QED**

Acknowledgements. The first author is very grateful to her advisor Professor P. T. Nagy for his useful comments during this work. The second author was supported by the grant GA ĆR 201/05/2707 and the project MSM 0021620839.

References

- [1] J. BERNDT, F. TRICCI, L. VANHECKE: Generalized Heisenberg Groups and Damek-Ricci Harmonic Spaces, Lecture Notes in Math., Vol. 1598, Springer-Verlag 1995.
- [2] P. EBERLEIN: *Geometry of 2-step nilpotent groups with a left invariant metric*, Ann. Sci. Ecole Norm. Sup., 4^e série, **27**, (1994), 611–660.
- [3] P. EBERLEIN: *Geometry of 2-step nilpotent groups with a left invariant metric II*, Trans. Amer. Math. Soc., **343**, (1994), 805–828.
- [4] R. GORNET, M. B. MAST: *The length spectrum of Riemannian two-step nilmanifolds*, Ann. Sci. Ecole Norm. Sup., 4^e série, **33**, (2000), 181–209.
- [5] S. HOMOLYA, P. T. NAGY: *Submersions on nilmanifolds and their geodesics*, Publ. Math. Debrecen, **62/3-4**, (2003), 415–428.
- [6] A. KAPLAN: *Riemannian nilmanifolds attached to Clifford modules*, Geometriae Dedicata, **11**, (1981), 127–136.
- [7] A. KAPLAN: *On the geometry of groups of Heisenberg type*, Bull. London Math. Soc., **15**, (1983), 35–42.
- [8] O. KOWALSKI, L. VANHECKE: *Classification of five-dimensional naturally reductive spaces*, Math. Proc. Camb. Phil. Soc., **97**, (1985), 445–463.
- [9] J. LAURET: *Homogeneous nilmanifolds of dimension 3 and 4*, Geometriae Dedicata, **68**, (1997), 145–155.
- [10] J. LAURET: *Modified H-type groups and symmetric-like Riemannian spaces*, Diff. Geom. Appl., **10**, (1999), 121–143.
- [11] L. MAGNIN: *Sur les algèbres de Lie nilpotents de dimension ≤ 7* , J. Geom. Phys., **3/1**, (1986), 119–144.
- [12] E. WILSON: *Isometry groups on homogeneous nilmanifolds*, Geometriae Dedicata, **12**, (1982), 337–346.